Author Index

Åkesson-Johansson, A., see Harmenberg, J. 75
Allender, W. J.
Gas–liquid chromatographic analysis of ethyl fenthion in tissues of native birds 199
Almeida, M. Lurdes, S., see Fransson, B. 229
Antus, S.
——, Bauer, R., Gottsegen, A. and Wagner, H.
Enantiomeric separation of racemic pterocarpans by high-performance liquid chromatography on (+)-poly(triphenylmethyl methacrylate)-coated silica gel 212
Baldonado, I. P., see Canova-Davis, E. 81
Bard, M., see Goss, T. A. 279
Bauer, R., see Antus, S. 212
Beckers, J. L.
——, and Everaerts, F. M.
Isotachophoresis with two leading ions and migration behaviour in capillary zone electrophoresis. I. Isotachophoresis with two leading ions 3
——, and Everaerts, F. M.
Isotachophoresis with two leading ions and migration behaviour in capillary zone electrophoresis. II. Migration behaviour in capillary zone electrophoresis 19
Bermejo, J., see Díez, M. A. 363
Berthod, A., see Campiglia, A. D. 37
Bicchi, C.
——, and Pisciotta, A.
Use of two-dimensional gas chromatography in the direct enantiomer separation of chiral essential oil components 341
Blanchardie, P., see Denis, M. 97
Blanco, C. G., see Díez, M. A. 363
Blank, M. L.
——, Cress, E. A., Fitzgerald, V. and Snyder, F.
Thin-layer and high-performance liquid chromatographic separation of glycerolipid subclasses as benzoates. Derivatives of ether and ester analogues of phosphatidylcholine, phosphatidylethanolamine and platelet activating factor 382
Brinkman, U. A. Th., see Veltkamp, A. C. 301
Burton, S. J.
——, Stead, C. V. and Lowe, C. R.
Design and applications of biomimetic anthraquinone dyes. III. Anthraquinone-immobilised C.I. Reactive Blue 2 analogues and their interaction with horse liver alcohol dehydrogenase and other adenine nucleotide-binding proteins 109
Camilleri, P.
——, Murphy, J. A. and Thorpe, C.
Direct chromatographic separation of the enantiomers of oxiracetam 208
Campiglia, A. D.
——, Berthod, A. and Winefordner, J. D.
Solid-surface room-temperature phosphorescence detection for high-performance liquid chromatography 37
Canova-Davis, E.
——, Baldonado, I. P. and Teshima, G. M.
Characterization of chemically synthesized human relaxin by high-performance liquid chromatography 81
Carlson, Å.
——, and Gyllenhaal, O.
Separation of carboxylic acid enantiomers by gas chromatography after rapid derivatization with (*R*)- or (*S*)-1-phenylethylamine after activation by ethyl chloroformate 333
Cavazzutti, G., see Gagliardi, L. 252
Claessens, H. A., see Sparidans, R. W. 319
Cox, R. P., see Juhler, R. K. 232
Cox, S., see Harmenberg, J. 75
Cress, E. A., see Blank, M. L. 382
Crothers, D. M., see Maa, Y.-F. 61
Das, H. A., see Veltkamp, A. C. 301
Deeken, R. A., see Pogány, S. A. 179
Denis, M.
——, Blanchardie, P., Orsonneau, J.-L. and Lustenberger, P.
One-step 10^4-fold purification of transformed glucocorticoid receptor. Method for purifying receptors associated with M_r *ca.* 90 000 heat-shock protein 97
Díez, M. A.
——, Guillén, M. D., Blanco, C. G. and Bermejo, J.
Chromatographic study of methylcyclopentadiene dimers and iso-dimers and determination of their boiling points 363
Doremaele, G. H. J. van, see Sparidans, R. W. 319
Eppert, G.
——, Liebscher, G. and Stief, C.
Examination of organic trace contamination and thermo-oxidative deterioration of ε-caprolactam by high-performance liquid chromatography 149
Everaerts, F. M., see Beckers, J. L. 3, 19
Fitzgerald, V., see Blank, M. L. 382
Frankenberger, Jr., W. T., see Mehra, H. C. 265

Fransson, B.
——, Lurdes S. Almeida, M., Grehn, L. and Ragnarsson, U.
Separation of spermidine derivatives by reversed-phase high-performance liquid chromatography using a UV-absorbing counter ion 229
Fujii, Y.
——, Ikeda, Y., Okamoto, I. and Yamazaki, M.
Separation and determination of lanatosides in *Digitalis lanata* leaves by high-performance liquid chromatography 241
Gagliardi, L.
——, Cavazzutti, G., Turchetto, L., Manna, F. and Tonelli, D.
Determination of preservatives in cosmetic products by reversed-phase high-performance liquid chromatography. IV 252
Gavlick, W. K.
——, Stuff, J. R. and Uden, P. C.
Comparison of discharge tubes for deuterium-specific gas chromatographic detection with a microwave-induced plasma 187
Golshan-Shirazi, S., see Lin, B. 27
Goss, T. A.
——, Bard, M. and Jarrett, H. W.
High-performance affinity chromatography of DNA 279
Gottsegen, A., see Antus, S. 212
Grehn, L., see Fransson, B. 229
Guillén, M. D., see Díez, M. A. 363
Guiochon, G., see Lin, B. 27
Gyllenhaal, O., see Carlson, Å. 333
Harmenberg, J.
——, Cox, S. and Åkesson-Johansson, A.
Improved sample preparation method for high-performance liquid chromatography of deoxyribonucleoside thriphosphates from cell culture extracts 75
Hasler, A.
——, Sticher, O. and Meier, B.
High-performance liquid chromatographic determination of five widespread flavonoid aglycones 236
Hayes, G. E.
Direct determination of corrosion inhibitor in aviation fuel by a column-switching technique using gel-permeation chromatography 259
Herk, A. M. van, see Sparidans, R. W. 319
Hirayama, N.
——, and Kuwamoto, T.
Influence of dissociation equilibria on the elution behaviour of the sample anion in anion chromatography 51
Ho, J. W.
Separation of porphyrins on cyclodextrin-bonded phases with a novel mobile phase 375
Hofmann, U.
——, Holzer, S. and Meese, C. O.
Pentafluorophenyldiazoalkanes as novel derivatization reagents for the determination of sensitive carboxylic acids by gas chromatography–negative-ion mass spectrometry 349
Holzer, S., see Hofmann, U. 349
Horie, M., see Saito, K. 141
Horváth, Cs., see Maa, Y.-F. 61
Hoshino, Y., see Saito, K. 141
Huisden, R.E.
——, Kraak, J. C. and Poppe, H.
Modification of silica with glucose for the separation of proteins by high-performance liquid chromatography 289
Huysmans, K. D., see Mehra, H. C. 265
Hynning, P.-Å., see Remberger, M. 159
Ikeda, Y., see Fujii, Y. 241
Inczédy, J.
—— and Szokoli, F.
Contributions to the calculation of retention data in ion-pair chromatography 309
Ivanova, L. A., see Karabonov, N. T. 193
Iwamoto, K., see Seno, M. 127
Jarrett, H. W., see Goss, T. A. 279
Jonhson, B. J.
Sulfite–disulfite equilibrium on an ion chromatography column 271
Juhler, R. K.
——, and Cox, R. P.
High-performance liquid chromatographic determination of chloroplast pigments with optimized separation of lutein and zeaxanthin 232
Kaniansky, D., see Kenndler, E. 203
Karabanov, N. T.
——, Vetrova, Z. P., Ivanova, L. A., Schuvalova, T. N. and Yashin, Ya. I.
Modified sorbents in binary stationary phase systems in gas chromatography 193
Katakawa, J., see Yasaka, Y. 133
Kenndler, E.
——, Schwer, C. and Kaniansky, D.
Purity control of riboflavin-5′-phosphate (vitamin B_2 phosphate) by capillary zone electrophoresis 203
Kikuchi, T., see Takatsuki, K. 357
Kobayashi, K., see Suzuki, S. 225
Kraak, J. C., see Huisden, R. E. 289
Kuwamoto, T., see Hirayama, N. 51
Liebscher, G., see Eppert, G. 149
Lin, B.
——, Golshan-Shirazi, S. and Guiochon, G.
Perturbation analysis of the influence of the isotherm coupling factor on the resolution between two compounds in chromatography 27

Lin, M., see Seno, M. 127
Lin, S.-C., see Maa, Y.-F. 61
Lowe, C. R., see Burton, S. J. 109
Lurdes S. Almeida, M., see Fransson, B. 229
Lustenberger, P., see Denis, M. 97
Maa, Y.-F.
——, Lin, S.-C., Horváth, Cs., Yang, U.-C. and Crothers, D. M.
Rapid high-performance liquid chromatography of nucleic acids with polystyrene-based micropellicular anion exchangers 61
MacNeil, J. D., see VanderKop, P. A. 386
Mahoney, N. E.
——, and Roitman, J. N.
High-performance liquid chromatographic analysis of phenylpyrroles produced by *Pseudomonas cepacia* 247
Manna, F., see Gagliardi, L. 252
Meese, C. O., see Hofmann, U. 349
Mehra, H. C.
——, Huysmans, K. D. and Frankenberger, Jr., W. T.
Determination of borate at trace levels in environmental samples by ion-exclusion chromatography 265
Meier, B., see Hasler, A. 236
Murphy, J. A., see Camilleri, P. 208
Nakazawa, H., see Okamoto, M. 217
——, see Saito, K. 141
Neilson, A. H., see Remberger, M. 159
Noda, J., see Suzuki, S. 225
Nose, N., see Saito, K. 141
Okamoto, I., see Fujii, Y. 241
Okamoto, M.
——, and Nakazawa, H.
Optical resolution of abscisic acid metabolites using an ovomucoid-conjugated high-performance liquid chromatographic column 217
Orsonneau, J.-L., see Denis, M. 97
Pisciotta, A., see Bicchi, C. 341
Pogány, S. A.
——, Deeken, R. A. and Zentner, G. M.
Gas chromatographic assay for 3,9-diethylene-2,4,8,10-tetraoxaspiro[5.5]undecane 179
Poppe, H., see Huisden, R. E. 289
Ragnarsson, U., see Fransson, B. 229
Remberger, M.
——, Hynning, P.-Å. and Neilson, A. H.
Gas chromatographic analysis and gas chromatographic–mass spectrometric identification of components in the cyclohexane-extractable fraction from contaminated sediment samples 159
Řezanka, T.
——, and Sokolov, M. Yu.
Identification of fatty acids longer than C_{32} in a sulphate-reducing bacterium by reversed-phase high-performance liquid chromatography and gas chromatography–mass spectrometry 275
Roitman, J. N., see Mahoney, N. E. 247
Saito, K.
——, Horie, M., Hoshino, Y., Nose, N. and Nakazawa, H.
High-performance liquid chromatographic determination of glycoalkaloids in potato products 141
Schuvalova, T. N., see Karabonov, N. T. 193
Schwer, C., see Kenndler, E. 203
Seno, M.
——, Lin, M. and Iwamoto, K.
Chromatographic behaviour of cyclodextrin complexes in NADH and NADP 127
Shono, T., see Yasaka, Y. 133
Snyder, F., see Blank, M. L. 382
Sokolov, M. Yu., see Řezanka, T. 275
Sparidans, R. W.
——, Claessens, H. A., Van Doremaele, G. H. J. and Van Herk, A. M.
Analysis of poly(styrene-co-methyl acrylate) and poly(styrene–co-butyl acrylate) by high-performance liquid chromatography 319
Stead, C. V., see Burton, S. J. 109
Sticher, O., see Hasler, A. 236
Stief, C., see Eppert, G. 149
Stobiecki, M.
——, and Wojtaszek, P.
Application of gas chromatography–mass spectrometry to the identification of isoflavonoids in lupine root extracts 391
Stuff, J. R., see Gavlick, W. K. 187
Sun, Z.-P., see Zhou, Y. 220
Suzuki, S.
——, Kobayashi, K., Noda, J., Suzuki, T. and Takama, K.
Simultaneous determination of biogenic amines by reversed-phase high-performance liquid chromatography 225
Suzuki, T., see Suzuki, S. 225
Szokoli, F., see Inczédy, J. 309
Takama, K., see Suzuki, S. 225
Takatsuki, K.
——, and Kikuchi, T.
Determination of N-nitrosodimethylamine in fish products using gas chromatography with nitrogen-phosphorus detection 357
Tanaka, M., see Yasaka, Y. 133
Teshima, G. M., see Canova-Davis, E. 81
Tetsumi, T., see Yasaka, Y. 133
Thorpe, C., see Camilleri, P. 208

Tonelli, D., see Gagliardi, L. 252
Turchetto, L., see Gagliardi, L. 252
Uden, P. C., see Gavlick, W. K. 187
VanderKop, P. A.
——, and MacNeil, J. D.
Separation and detection of monensin, lasalocid and salinomycin by thin-layer chromatography/bioautography 386
Van Doremaele, G. H. J., see Sparidans, R. W. 319
Van Herk, A. M., see Sparidans, R. W. 319
Veltkamp, A. C.
——, Das, H. A. and Brinkman, U. A. Th.
Direct and reverse flow-through heterogeneous scintillation counting of radiolabelled amino acids using post-column solvent segmentation 301
Vetrova, Z. P., see Karabonov, N. T. 193
Wagner, H., see Antus, S. 212
Winefordner, J. D., see Campiglia, A. D. 37
Wojtaszek, P., see Stobiecki, M. 391
Yamazaki, M., see Fujii, Y. 241
Yang, U.-C., see Maa, Y.-F. 61
Yasaka, Y.
——, Tanaka, M., Shono, T., Tetsumi, T. and Katakawa, J.
2-(2,3-Naphthalimino)ethyl trifluoromethanesulphonate as a highly reactive ultraviolet and fluorescent labelling agent for the liquid chromatographic determination of carboxylic acids 133
Yashin, Ya. I., see Karabonov, N. T. 193
Zentner, G. M., see Pogány, S. A. 179
Zhou, Y.
——, and Sun, Z.-P.
Liquid chromatographic evaluation of chiral derivatizing reagents for the resolution of amine and alcohol enantiomers 220

Erratum

J. Chromatogr., 504 (1990) 464–468

Page 466, Table I, last line of first column should read "$K_3[Co(Ox)_3]$" (where Ox = oxalate).